

Appl. No. 09/739,802

AMENDMENTS TO THE CLAIMS

1. (Canceled)
2. (Canceled)
3. (Canceled)
4. (Canceled)
5. (Canceled)
6. (Canceled)
7. (Canceled)

8. (Currently Amended) A catalyst for addition polymerization obtained by contacting:

(A) a compound containing an atom of the Group II to the Group XII or Lanthanide series of the Periodic Table of the Elements and containing no element of Group XIII, in which the lowest energy level of unoccupied molecular orbital having the valence p-type atomic orbital of the atom of the Group II to the Group XII or Lanthanide series as a main component wherein the coefficient represented by a linear combination is 0.4 or more is calculated to be 0.008 atomic unit (Hartree) or less by the calculation of density functional method (B3LYP/3-21G level) and wherein the compound (A) is a ~~perphyrin~~ or phthalocyanine complex in which a metal atom of the Group II to the Group XII or Lanthanide series is coordinated; τ with

Appl. No. 09/739,802

(B) a metal compound selected from the group consisting of compounds represented by ~~the general formula [4]:~~ $L_a M^1 X_b$ ~~[4]~~ and μ -oxo type compounds thereof, wherein M^1 is a metal atom of the Group III to the Group XIII or Lanthanide series; L is a group having cyclopentadienyl type anion skeleton or a group containing a hetero atom, a plurality of L's may be linked directly, or through a residual group containing a carbon atom, a silicon atom, a nitrogen atom, an oxygen atom, a sulfur atom or a phosphorous atom; X is a halogen atom or a hydrocarbon group; "a" represents a number satisfying $0 < a \leq 8$; and "b" represents a number satisfying $0 < b \leq 8$, ~~and μ -oxo type compounds thereof.~~

9. (Currently Amended) A catalyst for addition polymerization obtained by contacting:

(A) a compound containing an atom of the Group II to the Group XII or Lanthanide series of the Periodic Table of the Elements and containing no element of the Group XIII, in which the lowest energy level of unoccupied molecular orbital having the valence p-type atomic orbital of the atom of the Group II to the Group XII or Lanthanide series as a main component wherein the coefficient represented by a linear combination is 0.4 or more is calculated to be 0.008 atomic unit (Hartree) or less by the calculation of density functional method (B3LYP/3-21G level) and wherein the compound (A) is a ~~porphyrin or~~ phthalocyanine complex in which a

Appl. No. 09/739,802

metal atom of the Group II to the Group XII or Lanthanide series is coordinated; τ with


(B) a metal compound selected from the group consisting of compounds represented by the general formula ~~[4]~~: $L_a M^1 X_b$ ~~[4]~~ and μ -oxo type compounds thereof, wherein M^1 is a metal atom of the Group III to the Group XIII or Lanthanide series; L is a group having cyclopentadienyl type anion skeleton or a group containing a hetero atom, a plurality of L's may be linked directly, or through a residual group containing a carbon atom, a silicon atom, a nitrogen atom, an oxygen atom, a sulfur atom or a phosphorous atom; X is a halogen atom or a hydrocarbon group; "a" represents a number satisfying $0 < a \leq 8$; and "b" represents a number satisfying $0 < b \leq 8$, and ~~μ -oxo type compounds thereof~~, and an organoaluminum compound (C).

10. (Canceled)

11. (Canceled)

12. (Previously Amended) The catalyst according to claim 8, the compound (A) is a compound represented by the general formula [1]:


Appl. No. 09/739,802



wherein M represents an atom of the Group II to the Group XII or Lanthanide series of the Periodic Table, T represents an atom of the Group XIV or Group XV of the Periodic Table, and all of T's may be mutually the same or different, each of R¹ and R² independently is a hydrogen atom, a halogen atom, a hydrocarbon group or a halogenated hydrocarbon group, all of R¹'s and all of R²'s may be mutually the same or different, and may mutually form a ring, provided that at least one of R¹ and R² in the general formula [1] is an electron withdrawing group; X represents a hydrogen atom, a halogen atom, a hydrocarbon group or a hydrocarbon oxy group, and when a plural number of X's exist, they may be mutually the same or different, m represents a valence of M.

13. (Previously Amended) The catalyst according to claim 9, the compound (A) is a compound represented by the general formula [1]:

Appl. No. 09/739,802



wherein M represents an atom of the Group II to the Group XII or Lanthanide series of the Periodic Table, T represents an atom of the Group XIV or Group XV of the Periodic Table, and all of T's may be mutually the same or different, each of R¹ and R² independently is a hydrogen atom, a halogen atom, a hydrocarbon group or a halogenated hydrocarbon group, all of R¹'s and all of R²'s may be mutually the same or different, and may mutually form a ring, provided that at least one of R¹ and R² in the general formula [1] is an electron withdrawing group; X represents a hydrogen atom, a halogen atom, a hydrocarbon group or a hydrocarbon oxy group, and when a plural number of X's exist, they may be mutually the same or different, m represents a valence of M.

14. (Canceled)


15. (Canceled)

Appl. No. 09/739,802

16. (Previously Amended) The catalyst according to Claim 12, wherein the electron-withdrawing group is a fluorine, chlorine or bromine atom.

17. (Previously Amended) The catalyst according to Claim 13, wherein the electron-withdrawing group is a fluorine, chlorine or bromine atom.

18. (Previously Amended) The catalyst according to Claim 16, wherein the compound is a compound represented by the general formula [2]:




wherein M represents an atom of the Group II to the Group XII excluding Cu or Lanthanide series of the Periodic Table, X represents a hydrogen atom, a halogen atom, a hydrocarbon group or a hydrocarbon oxy group, and when a plural number of X's exist,

Appl. No. 09/739,802

they may be mutually the same or different, m represents a valence of M .

19. (Previously Amended) The catalyst according to Claim 17, wherein the compound is a compound represented by the general formula [2]:



wherein M represents an atom of the Group II to the Group XII excluding Cu or Lanthanide series of the Periodic Table, X represents a hydrogen atom, a halogen atom, a hydrocarbon group or a hydrocarbon oxy group, and when a plural number of X 's exist, they may be mutually the same or different, m represents a valence of M .

20. (Original) The catalyst according to Claims 18, wherein M is an atom of the Group IX or Group XII.

Appl. No. 09/739,802

21. (Original) The catalyst according to Claims 19, wherein M is an atom of the Group IX or Group XII.

22. (Original) The catalyst according to claim 8, wherein the compound(B) is a metallocene compound.

23. (Original) The catalyst according to claim 9, wherein the compound(B) is a metallocene compound.

24. (Withdrawn) A process for producing an addition polymer, which comprises polymerizing an addition polymerizable monomer in the presence of the catalyst of claim 8.

25. (Withdrawn) A process for producing an addition polymer, which comprises polymerizing an addition polymerizable monomer in the presence of the catalyst of claim 9.

26. (Withdrawn) A process for producing an addition polymer, which comprises polymerizing an addition polymerizable monomer in the presence of the catalyst of claim 22.

27. (Withdrawn) A process for producing an addition polymer, which comprises polymerizing an addition polymerizable monomer in the presence of the catalyst of claim 23.

Appl. No. 09/739,802

28. (Withdrawn) The process according to Claim 24, wherein the addition polymerizable polymer is an olefin.

29. (Withdrawn) The process according to Claim 25, wherein the addition polymerizable polymer is an olefin.

30. (Withdrawn) The process according to Claim 28, wherein the olefin is a mixture of ethylene and α -olefin.

31. (Withdrawn) The process according to Claim 29, wherein the olefin is a mixture of ethylene and α -olefin.